


# Inverse problems with one measurement

Eemeli Blåsten


Institute for Advanced Study,  
The Hong Kong University of Science and Technology

Inverse problems, PDE and geometry  
University of Jyväskylä  
Jyväskylä, August 22, 2018


# Scattering theory




=



+



## Mathematical scattering theory: measurements



Measurement:  $A_{u^i}$  is the **far-field pattern** of the scattered wave

$$u^s(x) = \frac{e^{ik|x|}}{|x|^{(n-1)/2}} A_{u^i} \left( \frac{x}{|x|} \right) + \mathcal{O} \left( \frac{1}{|x|^{n/2}} \right)$$

## Inverse problems

Given the far-field map  $u^j \mapsto A_{u^j}$ , recover  $V$  or its support  $\Omega$ .

# Inverse problems

Given the far-field map  $u^j \mapsto A_{u^j}$ , recover  $V$  or its support  $\Omega$ .

Solved when

- ▶ full far-field map given for all large frequencies (Saito 84),
- ▶ full far-field map given for a single frequency (Sylvester–Uhlmann 87  $n \geq 3$  + Bukhgeim 07  $n = 2$ ),
- ▶ + countless other variations

# Inverse problems

Given the far-field map  $u^j \mapsto A_{u^j}$ , recover  $V$  or its support  $\Omega$ .

Solved when



- ▶ full far-field map given for all large frequencies (Saito 84),
- ▶ full far-field map given for a single frequency (Sylvester–Uhlmann 87  $n \geq 3$  + Bukhgeim 07  $n = 2$ ),
- ▶ + countless other variations

Focus on *single measurement*:  $A_{u^j}$  given only for a single  $u^j$ .

Schiffer's problem: can a single measurement determine  $\Omega$ ?

# What about in physics?


Lord Rutherford's gold-foil experiment



Single incident wave

# Scattering theory

## Rutherford experiment's conclusions



measurement + *a-priori information* = conclusion

What if the measurement gives nothing?

# What if the measurement gives nothing?

Theorem (B.–Päivärinta–Sylvester CMP 14)

*The potential  $V = \chi_{[0, \infty[} \varphi$ ,  $\varphi(0) \neq 0$  always scatters.*

# What if the measurement gives nothing?

Theorem (B.–Päivärinta–Sylvester CMP 14)

*The potential  $V = \chi_{[0,\infty[} \varphi$ ,  $\varphi(0) \neq 0$  always scatters.*

For any incident wave  $u^i \neq 0$  we have  $A_{u^i} \neq 0$ .

## What if the measurement gives nothing?

Theorem (B.–Päivärinta–Sylvester CMP 14)

*The potential  $V = \chi_{[0,\infty[^n}\varphi$ ,  $\varphi(0) \neq 0$  always scatters.*

For any incident wave  $u^i \neq 0$  we have  $A_{u^i} \neq 0$ .

However  $A_{u^i}$  can become arbitrarily small with  $\|u^i\| \geq C > 0$ .  
(TE)

## Some follow-up corner scattering results

- ▶ Päivärinta–Salo–Vesalainen: 2D any angle, 3D almost any spherical cone
- ▶ Hu–Salo–Vesalainen: smoothness reduction, new arguments, *polygonal scatterer probing*
- ▶ Elschner–Hu: 3D any domain having two faces meet at an angle, and also curved edges
- ▶ Liu–Xiao: electromagnetic waves

## Some follow-up corner scattering results

- ▶ Päivärinta–Salo–Vesalainen: 2D any angle, 3D almost any spherical cone
- ▶ Hu–Salo–Vesalainen: smoothness reduction, new arguments, *polygonal scatterer probing*
- ▶ Elschner–Hu: 3D any domain having two faces meet at an angle, and also curved edges
- ▶ Liu–Xiao: electromagnetic waves

Injectivity of the Schiffer's problem for polyhedra:

### Theorem (HSV+EH)

Let  $P, P'$  be convex polyhedra and  $V = \chi_P \varphi$ ,  $V = \chi_{P'} \varphi'$  for admissible functions  $\varphi, \varphi'$ . Then

$$P \neq P' \implies A_{u^i} \neq A'_{u^i} \quad \forall u^i \neq 0$$

Any *single* incident wave determines  $P$  in the class of polyhedral penetrable scatterers.

## Some follow-up corner scattering results

- ▶ Päivärinta–Salo–Vesalainen: 2D any angle, 3D almost any spherical cone
- ▶ Hu–Salo–Vesalainen: smoothness reduction, new arguments, *polygonal scatterer probing*
- ▶ Elschner–Hu: 3D any domain having two faces meet at an angle, and also curved edges
- ▶ Liu–Xiao: electromagnetic waves

Injectivity of the Schiffer's problem for polyhedra:

### Theorem (HSV+EH)

Let  $P, P'$  be convex polyhedra and  $V = \chi_P \varphi$ ,  $V = \chi_{P'} \varphi'$  for admissible functions  $\varphi, \varphi'$ . Then

$$P \neq P' \implies A_{u^i} \neq A'_{u^i} \quad \forall u^i \neq 0$$

Any *single* incident wave determines  $P$  in the class of polyhedral penetrable scatterers.

However Ikehata's enclosure method gives roughly the same!

# My work while in Hong Kong

# Stability of polygonal scatterer probing

Non-vanishing total wave

Theorem (B., Liu, preprint)

Let  $u^i$  be an incident wave and let  $V = \chi_P \varphi$ ,  $V' = \chi_{P'} \varphi'$  be admissible with  $|u|, |u'| \neq 0$  in  $B_R$ . If

$$\|A_{u^i} - A'_{u^i}\|_{L^2(\mathbb{S}^{n-1})} < \varepsilon$$

then

$$d_H(P, P') \leq C(\ln \ln \|A_{u^i} - A'_{u^i}\|_2^{-1})^{-\eta}$$

for some  $\eta > 0$ .

# Lower bound for far-field pattern

Arbitrary Herglotz wave

Theorem (B., Liu, JFA 2017)

Let  $u^i$  be a normalized Herglotz wave,

$$u^i(x) = \int_{\mathbb{S}^{n-1}} e^{ik\theta \cdot x} g(\theta) d\sigma(\theta), \quad \|g\|_{L^2(\mathbb{S}^{n-1})} = 1,$$

and let  $V = \chi_P \varphi$  be admissible. Then

$$\|A_{u^i}\|_{L^2(\mathbb{S}^{n-1})} \geq C_{\|P_N\|, V} > 0$$

where the Taylor expansion of  $u^i$  at the corner  $x_c$  begins with  $P_N$ , and  $\|P_N\| = \int_{\mathbb{S}^{n-1}} |P_N(\theta)| d\sigma(\theta)$ .

## Mistake?



F. Cakoni: “Incident waves that approximate transmission eigenfunctions produce arbitrarily small far-field patterns.”

## From apparent contradiction to inspiration

Theorem (B., Liu, + B., Li, Liu, Wang, JFA + IP 2017 + preprint)

*Let the potential  $V = \chi_{\Omega}\varphi$  be admissible. Let  $v, w \neq 0$  be transmission eigenfunctions:*


$$\begin{aligned}(\Delta + k^2)v &= 0, & \Omega \\(\Delta + k^2(1 + V))w &= 0, & \Omega \\w - v &\in H_0^2(\Omega).\end{aligned}$$

Under  $C^\alpha$ -smoothness of  $v$  near  $x_c$ , we have

$$v(x_c) = w(x_c) = 0$$

at every corner point  $x_c$  of  $\Omega$ .

# Transmission eigenfunction localization



## Piecewise constant recovery

Injectivity of piecewise constant potential probing:

Theorem (B., Liu, preprint)

Let  $\Sigma_j, j = 1, 2, \dots$  be bounded convex polyhedra in an admissible geometric arrangement (think pixels/voxels) and  $V = \sum_j V_j \chi_{\Sigma_j}$ ,  $V = \sum_j V'_j \chi_{\Sigma_j}$  for constants  $V_j, V'_j \in \mathbb{C}$ . Then

$$V \neq V' \implies A_{u^i} \neq A'_{u^i} \quad \forall u^i(x) = e^{ik\theta \cdot x}$$

if  $k > 0$  small or  $|u| + |u'| \neq 0$  at each vertex.


A single incident plane wave determines  $V$  in the class of discretized penetrable scatterers.

## Generalizations and limitations

- ▶ unique determination of corner location *and* value


## Generalizations and limitations

- ▶ unique determination of corner location *and* value
- ▶ if  $\Sigma_j$  not known in advance: both  $(\Sigma_j)_{j=1}^{\infty}$  and  $V = \sum_j V_j \chi_{\Sigma_j}$  uniquely determined by a single measurement if geometry known to be **nested**





## Generalizations and limitations

- ▶ unique determination of corner location *and* value
- ▶ if  $\Sigma_j$  not known in advance: both  $(\Sigma_j)_{j=1}^{\infty}$  and  $V = \sum_j V_j \chi_{\Sigma_j}$  uniquely determined by a single measurement if geometry known to be **nested**



- ▶ method cannot yet be shown to distinguish between



# Non-scattering

Inverse source problem

$$(\Delta + k^2)u = f, \quad \lim_{r \rightarrow \infty} (\partial_r - ikr)u = 0$$

can one have  $f \neq 0$  but  $u_\infty = 0$ ?

# Non-scattering

## Inverse source problem

$$(\Delta + k^2)u = f, \quad \lim_{r \rightarrow \infty} (\partial_r - ikr)u = 0$$

can one have  $f \neq 0$  but  $u_\infty = 0$ ?

Recall:

$$u_\infty(\theta) = c_{k,n} \hat{f}(k\theta).$$

# Non-scattering

## Inverse source problem

$$(\Delta + k^2)u = f, \quad \lim_{r \rightarrow \infty} (\partial_r - ikr)u = 0$$

can one have  $f \neq 0$  but  $u_\infty = 0$ ?

Recall:

$$u_\infty(\theta) = c_{k,n} \hat{f}(k\theta).$$

Yes: let

$$f(x) = \begin{cases} 1, & |x| < r_0 \\ 0, & |x| \geq r_0 \end{cases}$$

where  $r_0 > 0$ .

# Non-scattering

## Inverse source problem

$$(\Delta + k^2)u = f, \quad \lim_{r \rightarrow \infty} (\partial_r - ikr)u = 0$$

can one have  $f \neq 0$  but  $u_\infty = 0$ ?

Recall:

$$u_\infty(\theta) = c_{k,n} \hat{f}(k\theta).$$

Yes: let

$$f(x) = \begin{cases} 1, & |x| < r_0 \\ 0, & |x| \geq r_0 \end{cases}$$

where  $r_0 > 0$ . Then

$$u_\infty(\theta) = c_{k,n} \hat{f}(k\theta) = c'_{k,n} J_{n/2}(kr_0) = 0$$

if  $kr_0$  is a zero of the Bessel function of order  $n/2$ .

# Always scattering

Smallness  $1/2$

A *small* uniform ball always scatters!

# Always scattering

## Smallness 1/2

A *small* uniform ball always scatters!

Also: any small scatterer always scatters!

### Theorem (B., Liu, preprint)

Let  $n \geq 2$ ,  $R_m, k \in \mathbb{R}_+$ ,  $0 \leq \alpha \leq 1$ . Let  $\Omega \subset \mathbb{R}^n$  be a bounded Lipschitz domain of diameter at most  $R_m$  and whose complement is connected. Let  $\Omega_c$  be a component of  $\Omega$ . The source  $f = \chi_{\Omega} \varphi$  radiates a non-zero far-field pattern at wavenumber  $k$  if

$$(\text{diam}(\Omega_c))^\alpha \leq C \frac{\sup_{\partial\Omega_c} |\varphi|}{\|\varphi\|_{C^\alpha(\bar{\Omega}_c)}},$$

for some  $C = C(k, R_m, n) > 0$ .

# Always scattering

Smallness 2/2

Proof.

Suppose  $(\Delta + k^2)u = \chi_{\Omega}\varphi$  and  $u_{\infty} = 0$ . Then  $u|_{\Omega^c} = 0$ , so  $u|_{\Omega_c} \in H_0^2(\Omega_c)$  and  $(\Delta + k^2)u = \varphi$  in  $\Omega_c$ .

# Always scattering

## Smallness 2/2

Proof.

Suppose  $(\Delta + k^2)u = \chi_{\Omega}\varphi$  and  $u_{\infty} = 0$ . Then  $u|_{\Omega^c} = 0$ , so

$u|_{\Omega_c} \in H_0^2(\Omega_c)$  and  $(\Delta + k^2)u = \varphi$  in  $\Omega_c$ .

Set  $g = \varphi - k^2u$ . Then elliptic regularity implies  $g \in C^\alpha(\bar{\Omega}_c)$  with  $\|g\|_{C^\alpha} \leq C(n, k, R_m) \|\varphi\|_{C^\alpha}$ . Moreover

$$\int_{\Omega_c} g(x) dx = \int_{\Omega_c} 1 \cdot \Delta u dx = 0$$

because  $u = \partial_\nu u = 0$  in  $\partial\Omega_c$ .

# Always scattering

## Smallness 2/2

Proof.

Suppose  $(\Delta + k^2)u = \chi_{\Omega}\varphi$  and  $u_{\infty} = 0$ . Then  $u|_{\Omega^c} = 0$ , so  $u|_{\Omega_c} \in H_0^2(\Omega_c)$  and  $(\Delta + k^2)u = \varphi$  in  $\Omega_c$ .

Set  $g = \varphi - k^2u$ . Then elliptic regularity implies  $g \in C^\alpha(\overline{\Omega_c})$  with  $\|g\|_{C^\alpha} \leq C(n, k, R_m) \|\varphi\|_{C^\alpha}$ . Moreover

$$\int_{\Omega_c} g(x) dx = \int_{\Omega_c} 1 \cdot \Delta u dx = 0$$

because  $u = \partial_\nu u = 0$  in  $\partial\Omega_c$ . Let  $p \in \partial\Omega_c$ . Then

$$\varphi(p)m(\Omega_c) = g(p)m(\Omega_c) = - \int_{\Omega_c} (g(x) - g(p)) dx$$

# Always scattering

## Smallness 2/2

Proof.

Suppose  $(\Delta + k^2)u = \chi_{\Omega}\varphi$  and  $u_{\infty} = 0$ . Then  $u|_{\Omega^c} = 0$ , so

$u|_{\Omega_c} \in H_0^2(\Omega_c)$  and  $(\Delta + k^2)u = \varphi$  in  $\Omega_c$ .

Set  $g = \varphi - k^2u$ . Then elliptic regularity implies  $g \in C^\alpha(\bar{\Omega}_c)$  with  $\|g\|_{C^\alpha} \leq C(n, k, R_m) \|\varphi\|_{C^\alpha}$ . Moreover

$$\int_{\Omega_c} g(x) dx = \int_{\Omega_c} 1 \cdot \Delta u dx = 0$$

because  $u = \partial_\nu u = 0$  in  $\partial\Omega_c$ . Let  $p \in \partial\Omega_c$ . Then

$$\varphi(p) m(\Omega_c) = g(p) m(\Omega_c) = - \int_{\Omega_c} (g(x) - g(p)) dx$$

Hence

$$|\varphi(p)| m(\Omega_c) \leq \|g\|_{C^\alpha} \int_{\Omega_c} |x - p|^\alpha dx \leq \|g\|_{C^\alpha} m(\Omega_c) (\text{diam}(\Omega_c))^\alpha.$$

# Always scattering

High curvature case

Is smallness the true cause for non-scattering?

# Always scattering

## High curvature case

Is smallness the true cause for non-scattering?

No: high curvature!

### Theorem (B., Liu, preprint)


*Let  $\Omega \subset \mathbb{R}^n$ ,  $n \geq 2$ , be a bounded domain. Assume that  $p \in \partial\Omega$  is an admissible  $K$ -curvature point. Also assume  $p$  connected to infinity outside of  $\Omega$ . Consider the source  $\chi_\Omega \varphi$ ,  $\varphi \in C^\alpha(\mathbb{R}^n)$ . If*

$$|\varphi(p)| \geq C(\ln K)^{(n+3)/2} K^{-\delta}$$

*then the source scatters a non-zero far-field pattern at wavenumber  $k$ .*

*Here  $\delta > 0$  depends on the geometric parameters, and  $C$  depends on the a-priori parameters of the  $K$ -curvature point, the wavenumber  $k$ , the upper bound for the diameter of  $\Omega$ , and the upper bound for  $\|\varphi\|_{C^\alpha}$ .*

## Admissible $K$ -curvature point



## Inverse source problem, Schiffer's problem

$$(\Delta + k^2)u = f = \chi_{\Omega}\varphi, \quad \lim_{r \rightarrow \infty} (\partial_r - ikr)u = 0$$

Can  $u_{\infty}(\theta) = \hat{c}\hat{f}(k\theta)$  determine  $\Omega$  given a fixed  $k$ ?

## Inverse source problem, Schiffer's problem

$$(\Delta + k^2)u = f = \chi_{\Omega}\varphi, \quad \lim_{r \rightarrow \infty} (\partial_r - ikr)u = 0$$

Can  $u_{\infty}(\theta) = \hat{c}f(k\theta)$  determine  $\Omega$  given a fixed  $k$ ?

Unique determination:

- ▶  $u_{\infty} = u'_{\infty} \implies \Omega = \Omega'$  for convex polyhedral shapes (corner scattering). Also for elasticity (with Lin), electromagnetism (with Liu, Xiao),
- ▶  $u_{\infty} = u'_{\infty} \implies \Omega \approx \Omega'$  for convex polyhedral shapes whose corners have been smoothed to admissible  $K$ -curvature points (high curvature scattering),
- ▶  $u_{\infty} = u'_{\infty} \implies \Omega \approx \Omega'$  for well-separated collections of small scatterers (small source scattering).

Thank you for your attention!