

Scattering from corners and other singularities

Emilia Blåsten

LUT University, Finland


Finnish Mathematical Days 2024

Aalto University

January 5, 2024

Scattering theory

Fixed frequency scattering



The total wave u satisfies


$$(\Delta + k^2(1 + V))u = 0,$$

V models a **perturbation** of the background,


$$u = u^i(x) + u^s(x)$$

\uparrow incident wave \leftarrow scattered wave


Scattering theory



=




+



$$u = u^i + u^s$$

Fixed frequency scattering theory: measurements



Measurement: A_{u^i} is the **far-field pattern** of the scattered wave

$$u^s(x) = \frac{e^{ik|x|}}{|x|^{(n-1)/2}} A_{u^i} \left(\frac{x}{|x|} \right) + \mathcal{O} \left(\frac{1}{|x|^{n/2}} \right)$$

Different inverse scattering problems

Given the **far-field map** $u^i \mapsto A_{u^i}$, recover the scattering potential V or its support Ω .

Different inverse scattering problems

Given the **far-field map** $u^i \mapsto A_{u^i}$, recover the scattering potential V or its support Ω .

Solved when

- ▶ full far-field map given for all large frequencies (Saito 1984)
- ▶ full far-field map given for a single frequency
 - ▷ Sylvester–Uhlmann 1987: 3D Calderón problem
 - ▷ R. Novikov 1988: 3D scattering
 - ▷ Bukhgeim 2007: 2D scattering
- ▶ + countless other variations

Different inverse scattering problems

Given the **far-field map** $u^j \mapsto A_{u^j}$, recover the scattering potential V or its support Ω .

Solved when

- ▶ full far-field map given for all large frequencies (Saito 1984)
- ▶ full far-field map given for a single frequency
 - ▷ Sylvester–Uhlmann 1987: 3D Calderón problem
 - ▷ R. Novikov 1988: 3D scattering
 - ▷ Bukhgeim 2007: 2D scattering
- ▶ + countless other variations

My focus is on **single measurement**: A_{u^j} given only for a single u^j .

Schiffer's problem: can a single measurement determine Ω ?


Why one measurement only?

Why one measurement only?

Example: Lord Rutherford's gold-foil experiment



Why one measurement only?

Example: Lord Rutherford's gold-foil experiment





Why one measurement only?

Example: Lord Rutherford's gold-foil experiment



Why one measurement only?


Example: Lord Rutherford's gold-foil experiment



Single incident wave

Scattering theory

Rutherford experiment's conclusions



measurement + a-priori information = conclusion

What if the measurement gives nothing?

It is very unfortunate if the far-field map is not injective. Most scattering potentials do have interior transmission eigenvalues. These exist when the map is non-injective. So it looks like the situation is unfortunate?

What if the measurement gives nothing?

It is very unfortunate if the far-field map is not injective. Most scattering potentials do have interior transmission eigenvalues. These exist when the map is non-injective. So it looks like the situation is unfortunate?

Theorem (B.–Päivärinta–Sylvester CMP 2014)

The potential $V = \chi_{[0,\infty[^n}\varphi$, $\varphi(0) \neq 0$ always scatters.

What if the measurement gives nothing?

It is very unfortunate if the far-field map is not injective. Most scattering potentials do have interior transmission eigenvalues. These exist when the map is non-injective. So it looks like the situation is unfortunate?

Theorem (B.-Päivärinta–Sylvester CMP 2014)

The potential $V = \chi_{[0,\infty[^n}\varphi$, $\varphi(0) \neq 0$ always scatters.

For **any** incident wave $u^i \neq 0$ and wavenumber $k > 0$ we have $A_{u^i} \neq 0$. The far-field map is injective despite there being transmission eigenvalues!

What if the measurement gives nothing?

It is very unfortunate if the far-field map is not injective. Most scattering potentials do have interior transmission eigenvalues. These exist when the map is non-injective. So it looks like the situation is unfortunate?

Theorem (B.–Päivärinta–Sylvester CMP 2014)

The potential $V = \chi_{[0,\infty[^n}\varphi$, $\varphi(0) \neq 0$ always scatters.

For **any** incident wave $u^i \neq 0$ and wavenumber $k > 0$ we have $A_{u^i} \neq 0$. The far-field map is injective despite there being transmission eigenvalues!

However, if k is a **transmission eigenvalue** A_{u^i} can become arbitrarily small with $\|u^i\| \geq 1$.

Proof sketch

Rellich's theorem and unique continuation imply $u = u^i$ in Ω^c so

$$k^2 \int V u^i u_0 dx = - \int_{\Omega} u_0 (\Delta + k^2(1 + V))(u - u^i) dx = 0$$

if $(\Delta + k^2(1 + V))u_0 = 0$ in Ω .

Proof sketch

Rellich's theorem and unique continuation imply $u = u^i$ in Ω^c so

$$k^2 \int V u^i u_0 dx = - \int_{\Omega} u_0 (\Delta + k^2(1 + V))(u - u^i) dx = 0$$

if $(\Delta + k^2(1 + V))u_0 = 0$ in Ω .

In simple case

$$u^i(x) = u^i(0) + u_r^i(x)$$

$$u_0(x) = e^{\rho \cdot x} (1 + \psi(x))$$

$$V(x) = \chi_{[0, \infty[}(\rho \cdot x) (\varphi(0) + \varphi_r(x))$$

Proof sketch

Rellich's theorem and unique continuation imply $u = u^i$ in Ω^c so

$$k^2 \int V u^i u_0 dx = - \int_{\Omega} u_0 (\Delta + k^2(1 + V))(u - u^i) dx = 0$$

if $(\Delta + k^2(1 + V))u_0 = 0$ in Ω .

In simple case

$$u^i(x) = u^i(0) + u_r^i(x)$$

$$u_0(x) = e^{\rho \cdot x} (1 + \psi(x))$$

$$V(x) = \chi_{[0, \infty[^n}(x) (\varphi(0) + \varphi_r(x))$$

Hölder estimates give

$$C |\varphi(0) u^i(0)| |\rho|^{-n} \leq \left| \varphi(0) u^i(0) \int_{[0, \infty[^n} e^{\rho \cdot x} dx \right| \leq C |\rho|^{-n-\delta}$$

if $\|\psi\|_p \leq C |\rho|^{-n/p-\varepsilon}$.

Some follow-up corner scattering results by others

- ▶ Päivärinta–Salo–Vesalainen 2017: 2D any angle, 3D almost any spherical cone
- ▶ Hu–Salo–Vesalainen 2016: smoothness reduction, new arguments, polygonal scatterer probing
- ▶ Elschner–Hu 2015, 2018: 3D any domain having two faces meet at an angle, and also curved edges
- ▶ Liu–Xiao 2017: electromagnetic waves
- ▶ ...
- ▶ free boundary methods:
 - ▷ Cakoni–Vogelius 2021: border singularities
 - ▷ Salo–Shahgholian 2021: analytic boundary non-scattering
 - ▷ ...

Lower bound for far-field pattern

Arbitrary Herglotz wave

Theorem (B.-Liu 2017)

Let u^i be a normalized Herglotz wave,

$$u^i(x) = \int_{\mathbb{S}^{n-1}} e^{ik\theta \cdot x} g(\theta) d\sigma(\theta), \quad \|g\|_{L^2(\mathbb{S}^{n-1})} = 1,$$

and let $V = \chi_P \varphi$ be admissible with x_c a corner of P . Then

$$\|A_{u^i}\|_{L^2(\mathbb{S}^{n-1})} \geq C_{\|P_N\|, V} > 0$$

where

$$\begin{aligned} u^i(x_c + r\theta) &= r^N P_N(\theta) + \mathcal{O}(r^{N+1}), \\ \|P_N\| &= \int_{\mathbb{S}^{n-1}} |P_N(\theta)| d\sigma(\theta) > 0. \end{aligned}$$

Mistake?



F. Cakoni: “Incident waves that approximate transmission eigenfunctions produce arbitrarily small far-field patterns.”

From apparent contradiction to inspiration

Theorem (B.-Liu 2017)

Let the potential $V = \chi_{\Omega}\varphi$ be admissible. Let $v, w \neq 0$ be transmission eigenfunctions:


$$\begin{aligned}(\Delta + k^2)v &= 0, & \Omega \\(\Delta + k^2(1 + V))w &= 0, & \Omega \\w - v &\in H_0^2(\Omega).\end{aligned}$$

Under C^α -smoothness of v near a convex corner x_c we have

$$v(x_c) = w(x_c) = 0.$$

Transmission eigenfunction localization

B.-Li-Liu-Wang 2017



Non-scattering

Technically simpler than potential scattering: inverse source problem

$$(\Delta + k^2)u = f, \quad \lim_{r \rightarrow \infty} (\partial_r - ikr) u = 0$$

Can one have $f \neq 0$ but $u_\infty = 0$?

Non-scattering

Technically simpler than potential scattering: inverse source problem

$$(\Delta + k^2)u = f, \quad \lim_{r \rightarrow \infty} (\partial_r - ikr) u = 0$$

Can one have $f \neq 0$ but $u_\infty = 0$?

Recall:

$$u_\infty(\theta) = c_{k,n} \hat{f}(k\theta).$$

I.e. can a compactly supported function have Fourier transform vanishing on a sphere?

Non-scattering

Technically simpler than potential scattering: inverse source problem

$$(\Delta + k^2)u = f, \quad \lim_{r \rightarrow \infty} (\partial_r - ikr)u = 0$$

Can one have $f \neq 0$ but $u_\infty = 0$?

Recall:

$$u_\infty(\theta) = c_{k,n} \hat{f}(k\theta).$$

I.e. can a compactly supported function have Fourier transform vanishing on a sphere?

Yes: let

$$f(x) = \begin{cases} 1, & |x| < r_0 \\ 0, & |x| \geq r_0 \end{cases}$$

where $r_0 > 0$.

Non-scattering

Technically simpler than potential scattering: inverse source problem

$$(\Delta + k^2)u = f, \quad \lim_{r \rightarrow \infty} (\partial_r - ikr)u = 0$$

Can one have $f \neq 0$ but $u_\infty = 0$?

Recall:

$$u_\infty(\theta) = c_{k,n} \hat{f}(k\theta).$$

I.e. can a compactly supported function have Fourier transform vanishing on a sphere?

Yes: let

$$f(x) = \begin{cases} 1, & |x| < r_0 \\ 0, & |x| \geq r_0 \end{cases}$$

where $r_0 > 0$. Then

$$u_\infty(\theta) = c_{k,n} \hat{f}(k\theta) = c'_{k,n} J_{n/2}(kr_0) = 0$$

if kr_0 is a zero of the Bessel function of order $n/2$.

Always scattering

Smallness $1/2$

A **small** uniform ball always scatters!

Always scattering

Smallness 1/2

A **small** uniform ball always scatters!

Also: any source with small shape always scatters!

Theorem (B.-Liu, 2021)

Let $n \geq 2$, $R_m, k \in \mathbb{R}_+$, $0 \leq \alpha \leq 1$. Let $\Omega \subset \mathbb{R}^n$ be a bounded Lipschitz domain of diameter at most R_m and whose complement is connected. Let Ω_c be a component of Ω . The source $f = \chi_{\Omega} \varphi$ radiates a non-zero far-field pattern at wavenumber k if

$$(\text{diam}(\Omega_c))^\alpha \leq C \frac{\sup_{\partial\Omega_c} |\varphi|}{\|\varphi\|_{C^\alpha(\overline{\Omega_c})}},$$

for some $C = C(k, R_m, n) > 0$.

Always scattering

Smallness 2/2: Proof

Suppose $(\Delta + k^2)u = \chi_{\Omega}\varphi$ and $u_{\infty} = 0$. Then $u|_{\Omega^c} = 0$, so $u|_{\Omega_c} \in H_0^2(\Omega_c)$ and $(\Delta + k^2)u = \varphi$ in Ω_c .

Always scattering

Smallness 2/2: Proof

Suppose $(\Delta + k^2)u = \chi_{\Omega}\varphi$ and $u_{\infty} = 0$. Then $u|_{\Omega^c} = 0$, so $u|_{\Omega_c} \in H_0^2(\Omega_c)$ and $(\Delta + k^2)u = \varphi$ in Ω_c .

Set $g = \varphi - k^2u$. Elliptic regularity implies $g \in C^\alpha(\overline{\Omega_c})$ with $\|g\|_\alpha \leq C(n, k, R_m) \|\varphi\|_\alpha$. Moreover $g = \Delta u$ and so

$$\int_{\Omega_c} g(x) dx = \int_{\Omega_c} 1 \cdot \Delta u dx = 0$$

because $u = \partial_\nu u = 0$ in $\partial\Omega_c$.

Always scattering

Smallness 2/2: Proof

Suppose $(\Delta + k^2)u = \chi_{\Omega}\varphi$ and $u_{\infty} = 0$. Then $u|_{\Omega^c} = 0$, so $u|_{\Omega_c} \in H_0^2(\Omega_c)$ and $(\Delta + k^2)u = \varphi$ in Ω_c .

Set $g = \varphi - k^2u$. Elliptic regularity implies $g \in C^\alpha(\overline{\Omega_c})$ with $\|g\|_\alpha \leq C(n, k, R_m) \|\varphi\|_\alpha$. Moreover $g = \Delta u$ and so

$$\int_{\Omega_c} g(x) dx = \int_{\Omega_c} 1 \cdot \Delta u dx = 0$$

because $u = \partial_\nu u = 0$ in $\partial\Omega_c$. Let $p \in \partial\Omega_c$. Then

$$\varphi(p)m(\Omega_c) = g(p)m(\Omega_c) = - \int_{\Omega_c} (g(x) - g(p)) dx$$

Always scattering

Smallness 2/2: Proof

Suppose $(\Delta + k^2)u = \chi_{\Omega}\varphi$ and $u_{\infty} = 0$. Then $u|_{\Omega^c} = 0$, so $u|_{\Omega_c} \in H_0^2(\Omega_c)$ and $(\Delta + k^2)u = \varphi$ in Ω_c .

Set $g = \varphi - k^2u$. Elliptic regularity implies $g \in C^\alpha(\overline{\Omega_c})$ with $\|g\|_\alpha \leq C(n, k, R_m) \|\varphi\|_\alpha$. Moreover $g = \Delta u$ and so

$$\int_{\Omega_c} g(x) dx = \int_{\Omega_c} 1 \cdot \Delta u dx = 0$$

because $u = \partial_\nu u = 0$ in $\partial\Omega_c$. Let $p \in \partial\Omega_c$. Then

$$\varphi(p) m(\Omega_c) = g(p) m(\Omega_c) = - \int_{\Omega_c} (g(x) - g(p)) dx$$

Hence

$$|\varphi(p)| m(\Omega_c) \leq \|g\|_\alpha \int_{\Omega_c} |x - p|^\alpha dx \leq \|g\|_\alpha m(\Omega_c) (\text{diam}(\Omega_c))^\alpha.$$

Inverse source problem, Schiffer's problem

$$(\Delta + k^2)u = f = \chi_{\Omega}\varphi, \quad \lim_{r \rightarrow \infty} (\partial_r - ikr)u = 0$$

Can $u_{\infty}(\theta) = c\hat{f}(k\theta)$ determine Ω when k is fixed?

Inverse source problem, Schiffer's problem

$$(\Delta + k^2)u = f = \chi_{\Omega}\varphi, \quad \lim_{r \rightarrow \infty} (\partial_r - ikr)u = 0$$

Can $u_{\infty}(\theta) = c\hat{f}(k\theta)$ determine Ω when k is fixed?

Unique determination:

- ▶ $u_{\infty} = u'_{\infty} \implies \Omega = \Omega'$ for convex polyhedral shapes (**corner scattering**). Assuming non-vanishing total waves, also for elasticity (B.-Lin 2018), electromagnetism (B.-Liu-Xiao 2021),
- ▶ $u_{\infty} = u'_{\infty} \implies \Omega \approx \Omega'$ for convex polyhedral shapes whose corners have been smoothed to admissible K -curvature points (**high curvature scattering**, B.-Liu 2021),
- ▶ $u_{\infty} = u'_{\infty} \implies \Omega \approx \Omega'$ for well-separated collections of small scatterers (**small source scattering**, B.-Liu 2021).

Non-spherical cones

Potential scattering

Let C be any cone whose cross-section K is star-shaped and $\chi_K \in H^\tau(\mathbb{R}^2)$ for some $\tau > 1/2$.

Theorem (B.–Pohjola 2022)

For any $\delta > 0$ there is a cone C_δ such that $d_H(C_\delta, C) < \delta$ and with the following property: potentials of the form

$$V = \chi_{C_\delta} \varphi$$

*where φ is smooth enough (roughly $C^{1/4}$) and non-zero at the vertex **always scatter**.*

Non-spherical cones

Source scattering (easier)

Theorem (B.–Pohjola 2022)

A source $f = \chi_C \varphi$ for $(\Delta + k^2)u = f$ scatters *for any* $k > 0$ when φ is smooth enough and non-zero at the vertex of the cone C when

$$\int_{\mathbb{S}^2 \cap C} Y_2^m dS \neq 0$$

for $m \in \{-2, -1, 0, +1, +2\}$ and Y_2^m is the spherical harmonic of degree 2. *This is true if C fits into a thin enough spherical cone.*

Non-spherical cones

Source scattering (easier)


Theorem (B.–Pohjola 2022)

A source $f = \chi_C \varphi$ for $(\Delta + k^2)u = f$ scatters *for any* $k > 0$ when φ is smooth enough and non-zero at the vertex of the cone C when

$$\int_{\mathbb{S}^2 \cap C} Y_2^m dS \neq 0$$

for $m \in \{-2, -1, 0, +1, +2\}$ and Y_2^m is the spherical harmonic of degree 2. *This is true if C fits into a thin enough spherical cone.*

“Thin enough” means $\cos \theta \leq 1/\sqrt{3}$. The magic angle is $\approx 54.74^\circ$.



Scattering screens

A **flat screen** $\Omega = \Omega_0 \times \{0\}$ with $\Omega_0 \subset \mathbb{R}^2$ simply connected, bounded and smooth. Scattering from such a screen:

$$\begin{aligned}(\Delta + k^2)u^s &= 0, & \mathbb{R}^3 \setminus \overline{\Omega}, \\ u^i + u^s &= 0, & \Omega, \\ r(\partial_r - ik)u^s &\rightarrow 0, & r = |x| \rightarrow \infty.\end{aligned}$$

Let Ω, Ω' be flat screens, $k > 0$, u^i an arbitrary incident wave, and $u^s, u^{s'}$ corresponding scattered waves.

Theorem (B.–Päivärinta–Sadique 2020)

- ▶ If $u^i(x_1, x_2, x_3) + u^i(x_1, x_2, -x_3) \neq 0$ for some x and $u_\infty^s = u_\infty^{s'}$ then $\Omega = \Omega'$.
- ▶ If $u^i(x_1, x_2, x_3) + u^i(x_1, x_2, -x_3) = 0$ for all x then $u_\infty^s = u_\infty^{s'} = 0$.

Thank you!